

*Pacific  
Journal of  
Mathematics*

**THE MAXIMAL SYSTOLE OF HYPERBOLIC SURFACES  
WITH MAXIMAL  $S^3$ -EXTENDABLE ABELIAN SYMMETRY**

YUE GAO AND JIAJUN WANG

Volume 325 No. 1

July 2023



# THE MAXIMAL SYSTOLE OF HYPERBOLIC SURFACES WITH MAXIMAL $S^3$ -EXTENDABLE ABELIAN SYMMETRY

YUE GAO AND JIAJUN WANG

**We study the maximal systole of hyperbolic surfaces with certain symmetries. We give the formula for the maximal systole of the surfaces that admit the largest  $S^3$ -extendable abelian group symmetry. The result is obtained by parametrizing such surfaces and enumerating all possible systoles.**

## 1. Introduction

The systole is an important topic in the study of hyperbolic surfaces. The systole has applications in various areas on surfaces, e.g, the Mumford’s classical compactness criterion [Mumford 1971] and the Weil–Peterson metric [Wolpert 2017; Wu 2019] in Teichmüller theory, and the spectrum of the Laplacian [Ballmann et al. 2016; 2018; Mondal 2014] and the optimal systolic ratio [Chen and Li 2015; Croke and Katz 2003; Gromov 1983] in differential geometry. For a survey on the study of the systole, see Parlier [2014].

We use the term “systole” to refer to either the minimal length of a closed geodesic on a hyperbolic surface, or a closed geodesic realizing this length, by abuse of notation. The systole can also be regarded as a real-valued function on the moduli space  $\mathcal{M}_g$  of all closed hyperbolic surfaces of genus  $g$  or the Teichmüller space  $\mathcal{T}_g$ . The maximal value of the systole function on  $\mathcal{M}_g$  is called the maximal systole in genus  $g$ . The maximal systole can be realized by Mumford’s compactness criterion. It is quite difficult to compute the exact value of the maximal systole. The only known case is genus 2, for which the maximal systole is realized by the Bolza surface [Jenni 1984].

It is also interesting to study the maximal value of the systole function on certain subspaces of  $\mathcal{M}_g$ . Bavard [1992] obtained the maximal systole of genera 2 and 5 on hyperelliptic surfaces. Schmutz [1993] gave a necessary and sufficient condition for the local maxima of the systole function and he gave some examples of local maxima with polyhedral symmetry. Fortier Bourque and Rafi [2022] constructed surfaces with locally maximal systoles and trivial symmetry.

Buser and Sarnak [1994] constructed surfaces with systoles larger than  $\frac{4}{3} \log g$  by arithmetic methods for infinitely many genera. Katz, Schaps, and Vishne [Katz

---

*MSC2020:* 30F45.

*Keywords:* systole, hyperbolic surfaces,  $S^3$ -extendable symmetry.

et al. 2007] obtained more surfaces with this lower bound. Hurwitz surfaces are among their examples. Petri and Walker [2018] and Petri [2018] gave concrete examples with systoles larger than  $\frac{4}{7} \log g - K$ .

Inspired by [Katz et al. 2007; Schmutz 1993], we are interested in the maximal systole of hyperbolic surfaces with certain symmetries. We consider hyperbolic surfaces with the largest  $S^3$ -extendable abelian symmetry. The  $S^3$ -extendable symmetry on a topological surface was introduced in [Wang et al. 2013; 2015]. A group action  $G$  on the genus- $g$  topological surface  $\Sigma_g$  is  $S^3$ -extendable if there exist an embedding  $i : \Sigma_g \rightarrow S^3$  and an injective homomorphism  $\phi : G \rightarrow \text{SO}(4)$  such that for any  $g \in G$ , the following diagram commutes:

$$\begin{array}{ccc} \Sigma_g & \xrightarrow{g} & \Sigma_g \\ \downarrow i & & \downarrow i \\ S^3 & \xrightarrow{\phi(g)} & S^3 \end{array}$$

When restricted to finite abelian groups, the maximal order of an  $S^3$ -extendable group action on  $\Sigma_g$  is  $2g + 2$  [Wang et al. 2013]. Such a group action can be realized as an isometry group action on a hyperbolic surface, and we say such a hyperbolic surface has the *maximal  $S^3$ -extendable abelian symmetry* or call the surface a *hyperbolic  $\Gamma(2, n)$  surface* [Wang et al. 2013] where  $n = g + 1$ .

Hyperbolic surfaces that admit an isometric  $S^3$ -extendable abelian group action of maximal order form a 2-dimensional subset of  $\mathcal{M}_g$ , and we consider the systole function on this subspace. Our main result is the following:

**Theorem 1.** *The maximal value of the systole function on hyperbolic  $\Gamma(2, n)$  surfaces is*

$$2 \operatorname{arccosh} K,$$

where

$$K = \sqrt[3]{\frac{1}{216}L^3 + \frac{1}{8}L^2 + \frac{5}{8}L - \frac{1}{8} + \sqrt{\frac{1}{108}L(L^2 + 18L + 27)}} \\ + \sqrt[3]{\frac{1}{216}L^3 + \frac{1}{8}L^2 + \frac{5}{8}L - \frac{1}{8} - \sqrt{\frac{1}{108}L(L^2 + 18L + 27)}} + \frac{1}{6}(L + 3),$$

and  $L = 4 \cos^2 \frac{\pi}{n}$ . The maximal value is obtained when

$$(c, t) = \left( \operatorname{arccosh} K, 2 \operatorname{arccosh} \frac{K+1}{2 \cos \frac{\pi}{n}} \right).$$

(The symbols  $c$  and  $t$  are defined in Section 2).

The maximal value of systoles of the  $\Gamma(2, n)$ -surfaces for small genera are shown in Table 1. We remark that the  $\Gamma(2, 3)$  surface is exactly the Bolza surface that realizes the maximal systole in genus 2.

genus	$\Gamma(2, n)$ maximal systole
2	3.0571
3	3.6478
4	3.9078
5	4.0464
6	4.1291

**Table 1.** Maximal systole of surface with largest  $S^3$ -extendable abelian symmetry.

Compared with the work [Bai et al. 2021] on the systole of surfaces with large cyclic symmetry for which the surfaces with large cyclic symmetry have a unique geometric structure, the surfaces studied in this paper form a two-dimensional subspace of  $\mathcal{M}_g$ , and the methods in the papers are quite different. We obtain our result by classifying the family of curves that are potentially the shortest geodesics on surfaces admitting this symmetry, and then determine when the systole is maximal.


The paper is organized as follows. In Section 2 we describe how to construct all hyperbolic  $\Gamma(2, n)$  surfaces and determine their symmetry. In Section 3, we give a useful lemma (Lemma 3) on the intersection properties of systoles. In Section 4, we prove that for any  $\Gamma(2, n)$  surface, there are only four closed geodesics in the quotient orbifold of the surface by its symmetric group that can lift to systoles of the  $\Gamma(2, n)$  surface (Proposition 7). In the last section, by calculating the length of these four curves and the differentials of these lengths, we get a condition for when the  $\Gamma(2, n)$  surface has the maximal systole (Proposition 9) and calculate its length.

## 2. The symmetry of $\Gamma(2, n)$ surfaces

In this section, we construct the hyperbolic  $\Gamma(2, n)$  surface and describe the geometry and topology of its quotient by its symmetry group.

Let  $\Sigma_{0,n}$  be the surface of genus 0 with  $n$  boundaries, endowed with a hyperbolic structure so that its boundaries are geodesics and  $\Sigma_{0,n}$  admits an isometric rotation of order  $n$ , as indicated in Figure 1. Each boundary circle is called a *cuff* of  $\Sigma$ . The shortest geodesic connecting two adjacent boundary circles or its image under the isometric rotation is called a *seam*. A seam is perpendicular to the two boundary circles it connects. The  $n$  seams cut  $\Sigma_{0,n}$  into two isometric right-angled hyperbolic  $2n$ -gons with geodesic boundary edges. The hyperbolic structure of  $\Sigma_{0,n}$  is parametrized by the length of its cuffs, called the *cuff length*. We denote the cuff length by  $2c$ , where  $c \in \mathbb{R}_+$  is called the *half cuff length*.

Two copies of  $\Sigma_{0,n}$  with the same cuff length can be glued together along the cuffs to form a closed surface, so that the two rotations on each copy can be extended




**Figure 1.**  $\Sigma_{0,n}$  and the right-angled  $2n$ -gon.

to the glued surface. We call the resulting surface a *hyperbolic  $\Gamma(2, n)$ -surface*. Two seams on the surface are *paired* if they connect the same two cuffs. Similar to the Fenchel–Nielsen coordinates on the Teichmüller space, a hyperbolic  $\Gamma(2, n)$ -surface can be parametrized by  $(c, t)$ , where  $c$  is the half cuff length and  $t$  is the “*twist parameter*”. The twist parameter  $t$  equals 0 if (any) two paired seams form a closed geodesic. The hyperbolic  $\Gamma(2, n)$  surface with parameter  $(c, t)$  is obtained from the hyperbolic  $\Gamma(2, n)$  surface with parameter  $(c, 0)$  by performing a Fenchel–Nielsen deformation of length  $t$  simultaneously along each cuff. Here the *Fenchel–Nielsen deformation* on a hyperbolic surface  $X$  along a simple closed geodesic  $\alpha \subset X$  with length  $t$  is constructed by cutting  $X$  along  $\alpha$  and then regluing the boundary curves with a left twist of length  $t$ . We may assume  $0 \leq t \leq c$  since the surface with parameter  $(c, t)$  is isometric to the surface with parameter  $(c, t + 2c)$  while the surface with parameter  $(c, t)$  is the reflection of the surface with parameter  $(c, 2c - t)$  when  $0 \leq t \leq 2c$ .

The symmetry group of a hyperbolic  $\Gamma(2, n)$  surface is  $D_n \oplus (\mathbb{Z}/2\mathbb{Z})$ , where  $D_n$  is the order  $n$  dihedral group. This symmetric group is generated by three rotations  $\sigma, \tau$ , and  $\rho$ . As illustrated in Figure 2,  $\sigma$  is the order  $n$  rotation that maps each  $n$ -holed sphere to itself,  $\tau$  is the order 2 rotation of each  $n$ -holed sphere, and  $\rho$  is the order 2 rotation exchanging the two  $n$ -holed spheres.

For a  $\Gamma(2, n)$  surface  $X$ , the quotient  $X/\langle \rho \rangle$  is a spherical orbifold with  $2n$  singular points of order 2, denoted as  $S^2(2, \dots, 2)_X$  (Figure 3, top left). The quotient  $X/\langle \rho, \sigma \rangle$  is a spherical orbifold with four singular points of order 2, 2,  $n, n$ , respectively, denoted as  $S^2(2, 2, n, n)_X$  (Figure 3, top right). The quotient  $X/\langle \rho, \sigma, \tau \rangle$  is a spherical orbifold with four singular points of order 2, 2, 2,  $n$ , respectively, denoted as  $S^2(2, 2, 2, n)_X$  (Figure 3, bottom). In Figure 3, top right,  $C$  and  $C'$  are the two order 2 singular points and  $O$  and  $O'$  are the order  $n$  singular points. In Figure 3, bottom,  $C, D, E$  are the order 2 singular points and  $O$  is the order  $n$  singular point. We will abbreviate the subscript  $X$  when there is no confusion. Denote the quotient




**Figure 2.** Generators for the isometry group of the hyperbolic  $\Gamma(2, n)$  surface.

branched covering maps as

$$(2-1) \quad X \begin{array}{c} \xrightarrow{p} \\ \xrightarrow{\pi} S^2(2, 2, \dots, 2)_X \xrightarrow{\pi'} S^2(2, 2, n, n)_X \xrightarrow{\pi''} S^2(2, 2, 2, n)_X \\ \xrightarrow{q} \end{array}$$

For a cuff  $\gamma_i$  in  $X$ , let  $A$  and  $\rho(A)$  be the endpoints of two paired seams between  $\gamma_i$  and  $\gamma_{i-1}$  on the cuff  $\gamma_i$ . The Fenchel–Nielsen deformation gives a geodesic of length  $t$  between  $A$  and  $\rho(A)$ , and we let  $C$  be the midpoint. The other two paired



**Figure 3.** Orbifolds for the hyperbolic  $\Gamma(2, n)$  surface.

seams give another midpoint  $C'$ . Then  $C$  and  $C'$  are fixed points of the rotation  $\rho$  on  $\gamma_i$  and  $|AC|$  is half the twist parameter  $t$ . It follows that in  $S^2(2, 2, n, n)_X$  (Figure 3, top right), we have

$$|CC'| = c, \quad |AC| = \frac{1}{2}t$$

and in  $S^2(2, 2, 2, n)_X$  (Figure 3, bottom) we have

$$|CE| = \frac{1}{2}c, \quad |AC| = \frac{1}{2}t.$$

### 3. The intersection properties of the systoles

In this section, we rule out curves on  $\Gamma(2, n)$  surface that cannot be the systole. The following lemma is classical and well known.

**Lemma 2.** *Any systole on a closed hyperbolic surface is simple and any two systoles intersect at most once. On the orbifold  $S^2(2, 2, \dots, 2)$ ,  $S^2(2, 2, n, n)$  or  $S^2(2, 2, 2, n)$ , any simple closed curve is separating, and any two simple closed curves are either disjoint or intersect at least twice.*

The following lemma is used to rule out curves on  $\Gamma(2, n)$  that cannot be the systole.


**Lemma 3.** *Given a hyperbolic  $\Gamma(2, n)$  surface  $X$ , let  $\pi, \pi', \pi''$  denote the branched covering maps*

$$X \xrightarrow{\pi} S^2(2, 2, \dots, 2)_X \xrightarrow{\pi'} S^2(2, 2, n, n)_X \xrightarrow{\pi''} S^2(2, 2, 2, n)_X$$

*Then under the maps  $\pi, \pi' \circ \pi$  and  $\pi'' \circ \pi' \circ \pi$ , the image of a systole on  $X$  has no self-intersection at any regular point on the targeting orbifold, and the images of two systoles do not intersect at any regular point on the targeting orbifold.*


*Proof.* (1) If  $\alpha$  is a simple closed curve in  $X$ ,  $\pi(\alpha)$  has a self-intersection point  $p$ . Then  $\pi^{-1}(p)$  consists of two points, both are the intersection points of  $\pi^{-1}(\pi(\alpha))$ . By the definition of double branched cover,  $\pi^{-1}(\pi(\alpha))$  consists of either one curve or two curves with equal length. Since  $\alpha$  is simple,  $\pi^{-1}(\pi(\alpha))$  consists of two curves. These two curves intersect at least twice, therefore cannot be systole.

We assume  $\alpha$  and  $\beta$  are two simple closed curves with equal length on  $X$ , Hence the shape of  $\pi(\alpha)$  and  $\pi(\beta)$  has two possibilities:  $S^1$  or a segment whose endpoints



**Figure 4.** Case (a):  $\pi(\alpha) \cup \pi(\beta)$ .





**Figure 5.** Case (a): The double covers of  $\pi(\alpha) \cup \pi(\beta)$ . The three types are shown in the top, the bottom left, and the bottom right, respectively.

are branched points of the branched cover  $\pi$ . We also assume  $p$  is the intersection point of  $\pi(\alpha)$  and  $\pi(\beta)$ .

(a) If  $\pi(\alpha)$  and  $\pi(\beta)$  are simple closed curves, then  $\pi(\alpha)$  intersects  $\pi(\beta)$  at least twice by Lemma 2. Recall that there are two types of double covers of  $S^1$ , namely  $S^1$  and  $S^1 \amalg S^1$ . Then there are three types of double covers of  $\pi(\alpha) \cup \pi(\beta)$ , shown in Figure 5.


In all the cases,  $\alpha$  intersects  $\beta$  at least twice, which contradicts to Lemma 2.

(b) If  $\pi(\alpha)$  is a segment while  $\pi(\beta)$  is a simple closed curve (Figure 6, left), then there are two types of the double (branched) covers of  $\pi(\alpha) \cup \pi(\beta)$  shown in Figure 6, middle and right.


If the double branched cover of  $\pi(\alpha) \cup \pi(\beta)$  is the case shown in Figure 6, middle, then it is clear that the curve  $\alpha$  and  $\beta$  have at least two intersections. Therefore,  $\alpha$  and  $\beta$  cannot be systoles.

If the double branched cover of  $\pi(\alpha) \cup \pi(\beta)$  is the case shown in Figure 6, right, we assume  $\tilde{p}$  is one of the branched point of  $\pi$  in Figure 6, right. Therefore  $\pi_*([\tilde{\beta}]) = \pi_*([\tilde{\beta}'])$  in  $\pi_1(\pi(X), \pi(\tilde{p}))$ . Here  $[\tilde{\beta}]$  and  $[\tilde{\beta}']$  are elements of  $\pi_1(X, \tilde{p})$  represented by  $\tilde{\beta}$  and  $\tilde{\beta}'$ . It contradicts the injectivity of  $\pi_*$  ( $\pi$  is a covering map).

(c) If both  $\pi(\alpha)$  and  $\pi(\beta)$  are segments (Figure 7), then  $|\pi^{-1}(\pi(\alpha)) \cap \pi^{-1}(\pi(\beta))| \geq 2$  since the intersection point of  $\pi(\alpha)$  and  $\pi(\beta)$  is a regular point. However, both  $\pi^{-1}(\pi(\alpha))$  and  $\pi^{-1}(\pi(\beta))$  are connected. Therefore  $|\alpha \cap \beta| \geq 2$ , so that  $\alpha$  and  $\beta$  cannot be systole.



**Figure 6.** Case (b):  $\pi(\alpha) \cup \pi(\beta)$  (left) and the double covers of  $\pi(\alpha) \cup \pi(\beta)$  (middle and right).



**Figure 7.** Case (c):  $\pi(\alpha) \cup \pi(\beta)$ .

(2) Let  $\alpha$  be a systole of  $X$ ; then by (1),  $\pi(\alpha)$  has no self-intersection and won't intersect the image of another systole at regular points. Therefore, if  $\pi'\pi(\alpha)$  has self-intersection at regular points, then it implies that either  $\pi(\alpha)$  intersects itself or it intersects another lift of  $\pi'\pi(\alpha)$ . Therefore  $\pi' \circ \pi(\alpha)$  has no self-intersections.

By exactly the same argument, we can prove that the images of two systoles of  $X$  on  $S^2(2, 2, n, n)$  do not intersect at any regular point of the orbifold.

The case for  $\pi'' \circ \pi' \circ \pi$  is similar to the case for  $\pi' \circ \pi$ . □

#### 4. The image of systoles on $S^2(2, 2, 2, n)$

For a  $\Gamma(2, n)$  surface  $X$ , we find geodesics in  $S^2(2, 2, 2, n)_X$  that lift to the systoles in  $X$  in this section.

**Lemma 4.** *For a  $\Gamma(2, n)$  surface  $X$ , a systole's image in the orbifold  $S^2(2, 2, 2, n)_X$  (Figure 3, bottom) has only two possibilities:*

- (1) A geodesic segment joining two order-two singular points ( $C$  and  $D$ ,  $C$  and  $E$  or  $D$  and  $E$ ).
- (2) A simple closed geodesic passing through  $C$ .


*Proof.* By Lemma 3, the image of a systole of  $X$  is a simple closed geodesic or a geodesic segment joining two singular points.

(1) Image of a systole of  $X$  cannot be a simple closed curve not passing through any singular point of the orbifold. Such a curve separates  $S^2(2, 2, 2, n)_X$  by Lemma 2. On each side of the curve, there are two singular points; otherwise, the curve lifts to null-homotopic curves in  $X$ . The order of both singular points on one side is two; hence the geodesic homotopic to this curve is the geodesic joining these two points.

(2) No systole's image passes through the order  $n$  singular point  $O$ . This point lifts to a regular point in  $S^2(2, 2, \dots, 2)_X$ , and a segment through  $O$  lifts to  $n$  segments intersecting at the preimage of  $O$ . Then by Lemma 3, this conclusion holds.

(3) The simple closed curve passing through  $D$  or  $E$  cannot lift to a systole of  $X$ , since  $D$  and  $E$  lift to regular points in  $S^2(2, 2, n, n)_X$ , and such curves lift to nonsimple curves (Figure 8).

By (1), (2), (3), this lemma holds. □



**Figure 8.** A simple closed curve passing through  $D$  in  $S^2(2, 2, 2, n)$  and its lift.

In order to obtain the systole of a  $\Gamma(2, n)$  surface, the next step is to find the geodesic in  $S^2(2, 2, 2, n)_X$  joining  $C$  and  $D$  (or  $C$  and  $E$  or  $D$  and  $E$  or the simple closed geodesic through  $C$ ), whose lift in  $X$  is the shortest one among all geodesics joining  $C$  and  $D$  (or  $C$  and  $E$  or  $D$  and  $E$  or the simple closed geodesic through  $C$ , respectively).


**Lemma 5.** For a  $\Gamma(2, n)$  surface  $X$ , let  $l$  and  $l'$  be two geodesics in  $S^2(2, 2, 2, n)_X$  joining  $C$  and  $D$  (or joining  $C$  and  $E$  or  $D$  and  $E$  or the simple closed geodesics passing through  $C$ ), and  $\tilde{l}$  and  $\tilde{l}'$  are their preimages in  $X$ , respectively. Then the covering  $\tilde{l} \rightarrow l$  and  $\tilde{l}' \rightarrow l'$  are topologically equivalent. More precisely, a homeomorphism  $f : l \rightarrow l'$  can lift to  $\tilde{f} : \tilde{l} \rightarrow \tilde{l}'$ , letting this diagram commute:

$$\begin{array}{ccc}
 \tilde{l} & \xrightarrow{\tilde{f}} & \tilde{l}' \\
 q \downarrow & & \downarrow q \\
 l & \xrightarrow{f} & l'
 \end{array}$$

*Proof.* We provide the proof for the geodesics joining  $C$  and  $D$  only, since the proofs for other cases are exactly the same.

For any  $l \subset S^2(2, 2, 2, n)_X$  joining  $C$  and  $D$ , there is a curve  $l''$  joining  $D$  and  $E$ , intersecting  $l$  only at  $D$ . The double branched cover  $\pi'' : S^2(2, 2, 2, n)_X \rightarrow S^2(2, 2, 2, n)_X$  can be constructed by gluing two copies of  $S^2(2, 2, 2, n)_X \setminus l''$  along their boundaries. Hence the preimage of  $l$  in  $S^2(2, 2, 2, n)_X$  is a segment (denoted as  $\tilde{l}_1$ ) joining  $C$  and  $C'$  for any  $l$ . Therefore the coverings of any two segments joining  $C$  and  $D$  are equivalent (Figure 9).

Similarly, for any  $\tilde{l}_1 \subset S^2(2, 2, 2, n)_X$  joining  $C$  and  $C'$ , there is a segment  $l''_1$  joining  $O$  and  $O'$ , intersecting  $\tilde{l}_1$  at exactly one point. Thus we can construct the  $n$ -fold cyclic branched cover of  $S^2(2, 2, 2, n)_X$  by gluing  $n$ -copies of  $S^2(2, 2, 2, n)_X \setminus l''_1$ . Since for any  $\tilde{l}_1$ , we always choose a curve  $l''_1$  intersecting  $\tilde{l}_1$  once, the covering



**Figure 9.**  $l$  and its lifts.

of  $\tilde{l}_1$  by its preimage in  $S^2(2, 2, \dots, 2)_X$  (denoted by  $\tilde{l}_2$ ) is topologically unique (Figure 9).

The multicurve  $\tilde{l}_2 \subset S^2(2, 2, \dots, 2)_X$  consists of segments joining the singular points. Therefore its preimage in the  $\Gamma(2, n)$  surface  $X$  (a manifold with no singular points) is topologically unique.  $\square$


**Corollary 6.** *Let  $l, l' \subset S^2(2, 2, 2, n)_X$  be geodesic segments joining  $C$  and  $D$ , and  $\alpha, \alpha' \subset X$  be simple closed geodesics lifted from  $l$  and  $l'$ , respectively. If  $|l| < |l'|$ , then  $|\alpha| < |\alpha'|$ .*

*This conclusion also holds for geodesics joining  $C$  and  $E$ , geodesics joining  $D$  and  $E$ , or simple closed geodesics passing through  $C$ .*


*Proof.* By Lemma 5,

$$\frac{|\alpha|}{|l|} = \frac{|\alpha'|}{|l'|}. \quad \square$$

**Proposition 7.** *For a  $\Gamma(2, n)$  surface  $X$ , there are only four possible geodesics in  $S^2(2, 2, 2, n)_X$  that lift to systoles in  $X$ . They are the shortest geodesics joining  $C$  and  $D$ , joining  $C$  and  $E$  and joining  $D$  and  $E$ , and the shortest simple closed geodesic passing through  $C$ , denoted as  $l_{CD}, l_{CE}, l_{DE}$  and  $l_C$ , respectively. Figure 10 describes the geometry of these four curves.*




**Figure 10.** The four geodesics that possibly lift to systoles of  $X$ .




**Figure 11.** Looking for the shortest geodesic joining  $C$  and  $D$  (top) and looking for the shortest simple closed geodesic passing through  $C$  (bottom).

*Proof.* By Corollary 6, the goal of this proposition is to describe the shortest geodesics in  $S^2(2, 2, 2, n)_X$  joining  $C$  and  $D$ , joining  $C$  and  $E$ , joining  $D$  and  $E$  and the simple closed geodesic passing through  $C$ .

(1) The shortest geodesic joining  $C$  and  $D$ : Let's consider the pentagon shown in Figure 11, a fundamental domain of  $S^2(2, 2, 2, n)_X$ . If a geodesic  $l \subset S^2(2, 2, 2, n)_X$  joining  $C$  and  $D$  consists of more than one segment in the pentagon (Figure 11, top left), then by reflecting some of its segments, we obtain a bending geodesic segment joining  $C$  and  $D$  or  $C$  and  $D_1$  with equal length to  $l$  and show that  $l$  is longer than the segment  $CD$  or  $CD_1$ .



**Figure 12.** Looking for the shortest geodesic joining  $C$  and  $D$  (I) (top) and looking for the shortest geodesic joining  $C$  and  $D$  (II) (bottom).



**Figure 13.** Cut and paste.

The segment  $CD$  is shorter than  $CD_1$ , because

$$|AD| = |AD_1| \quad \text{and} \quad |AC| < \frac{c}{2} < |A_1C|,$$

and it follows by the hyperbolic cosine law [Buser 2010, p. 454, 2.2.2 (i)]. Therefore, the shortest geodesic joining  $C$  and  $D$  is the segment  $l_{CD}$  shown in Figure 10, top left.

(2) The shortest geodesic joining  $D$  and  $E$  is the segment  $l_{DE}$  shown in Figure 10, top right, by the same argument.

(3) The shortest simple closed geodesic passing through  $C$  is the geodesic  $l_C$  shown in Figure 10, bottom right, by the same argument; see Figure 11.

(4) The shortest geodesic joining  $C$  and  $E$ : By reflecting some segments, we get the shortest geodesic is either the geodesic in Figure 12, top right, (denoted by  $l_{CE}$ ) or the geodesic in Figure 12, bottom right (denoted by  $l'_{CE}$ ). By the cut-and-paste shown in Figure 13, we see that  $l_{CE}$  is shorter than  $l'_{CE}$ , hence the shortest geodesic joining  $C$  and  $E$  is  $l_{CE}$  in Figure 10, bottom left.  $\square$

### 5. Calculations


In this section, we represent the length of the curves in Figure 10 by the parameters  $c$  and  $t$ . Then by these formulae, we give a condition of the  $\Gamma(2, n)$  surface having the longest systole. For convenience, we call this surface a *maximal surface*. Finally, we calculate the systole length of this surface.

Recall that in the pentagons in Figure 10,  $|CE| = c/2$ ,  $|AC| = t/2$ ,  $|A_1E| = (c - t)/2$  and  $\angle DOD_1 = 2\pi/n$ . We assume  $AD = A_1D_1 = s/2$ . Then in one of the two half pieces of the pentagon (Figure 14), by hyperbolic trigonometry [Buser 2010, p. 454, 2.3.1 (i)],

$$(5-1) \quad \sinh \frac{c}{2} \sinh \frac{s}{2} = \cos \frac{\pi}{n}.$$

Therefore, directly, for the lengths of the geodesics in Figure 10, we have

$$(5-2) \quad |l_{CE}| = \frac{c}{2},$$



**Figure 14.** The pentagon.

and by the hyperbolic cosine law in right-angled triangles [Buser 2010, p. 454, 2.2.2 (i)]

$$(5-3) \quad \cosh|l_{CD}| = \cosh|CD| = \cosh \frac{t}{2} \cosh \frac{s}{2};$$

$$(5-4) \quad \cosh|l_{DE}| = \cosh|D_1E| = \cosh \frac{c-t}{2} \cosh \frac{s}{2}.$$

To calculate the length of  $l_C$  (Figure 10, bottom right), we treat the pentagon as a fundamental domain for the orbifold  $S^2(2, 2, 2, n)$  in  $\mathbb{H}^2$ . Then in the joining of two pentagons shown in Figure 15, left,  $l_C$  is realized by the segment  $CC'$ . Hence its length is

$$(5-5) \quad \begin{aligned} \cosh|l_C| &= \cosh|CC'| \\ &= \cosh|AA_1| \cosh|AC| \cosh|A_1C'| - \sinh|AC| \sinh|A_1C'| \\ &= \cosh s \cosh \frac{t}{2} \cosh\left(c - \frac{t}{2}\right) - \sinh \frac{t}{2} \sinh\left(c - \frac{t}{2}\right) \end{aligned}$$

by a trigonometric formula [Buser 2010, p. 38, 2.3.2].


Now we are ready to prove the following:

**Proposition 8.** *In the  $\Gamma(2, n)$  surface  $X_0$  with maximal systole among all the  $\Gamma(2, n)$  surfaces,  $l_{DE}$  in  $S^2(2, 2, 2, n)_{X_0}$  cannot lift to a systole of this surface.*

*Proof.* Recall that by Proposition 7, in  $S^2(2, 2, 2, n)_{X_0}$ , there are only four geodesics that can lift to systoles of  $X_0$ , namely  $l_{CD}$ ,  $l_{DE}$ ,  $l_{CE}$  and  $l_C$  in Figure 10.

If  $l_{DE}$  lifts to a systole, then  $l_{CD}$  and  $l_{CE}$  cannot lift to systoles. This is because, when lifting to  $S^2(2, 2, n, n)$ ,  $l_{DE}$  intersects  $l_{CD}$  and  $l_{CE}$  at regular points  $D$  and  $E$  of the orbifold (Figure 15, right), therefore by Lemma 3, they cannot simultaneously become systoles.

Then we calculate the differentials of  $|l_{DE}|$  and  $|l_C|$ , showing that there is vector  $(A(c, t), B(c, t))$  such that  $d|l_C|(A, B) > 0$  and  $d|l_{DE}|(A, B) > 0$  simultaneously. Since only  $l_{DE}$  and  $l_C$  can lift to systoles, a surface with a systole lifted from  $l_{DE}$  cannot be a maximal surface.



**Figure 15.** The geodesic  $l_C$  (left) and the lift of  $l_{DE}$ ,  $l_{CD}$  and  $l_{CE}$  (right).

As a preparation, we differentiate both sides of (5-1) and get

$$(5-6) \quad \frac{ds}{dc} = -\frac{\cosh \frac{c}{2} \sinh \frac{s}{2}}{\cosh \frac{s}{2} \sinh \frac{c}{2}}.$$

Then for  $l_{DE}$

$$\begin{aligned} \frac{\partial |l_{DE}|}{\partial t} &= \frac{\partial}{\partial t} \left( \cosh \frac{s}{2} \cosh \frac{c-t}{2} \right) = -\frac{1}{2} \cosh \frac{s}{2} \sinh \frac{c-t}{2}, \\ \frac{\partial |l_{DE}|}{\partial c} &= \frac{\partial}{\partial c} \left( \cosh \frac{s}{2} \cosh \frac{c-t}{2} \right) \\ &= \frac{1}{2} \left( \sinh \frac{s}{2} \frac{ds}{dc} \cosh \frac{c-t}{2} + \cosh \frac{s}{2} \sinh \frac{c-t}{2} \right), \\ d|l_{DE}| &= \frac{\partial |l_{DE}|}{\partial t} dt + \frac{\partial |l_{DE}|}{\partial c} dc. \end{aligned}$$

For  $l_C$

$$(5-7) \quad \begin{aligned} \frac{\partial |l_C|}{\partial t} &= \frac{\partial}{\partial t} \left( \cosh s \cosh \frac{t}{2} \cosh \left( c - \frac{t}{2} \right) - \sinh \frac{t}{2} \sinh \left( c - \frac{t}{2} \right) \right) \\ &= \frac{1}{2} (\cosh s + 1) \sinh(t - c), \end{aligned}$$

and

$$\begin{aligned} \frac{\partial |l_C|}{\partial c} &= \frac{\partial}{\partial c} \left( \cosh s \cosh \frac{t}{2} \cosh \left( c - \frac{t}{2} \right) - \sinh \frac{t}{2} \sinh \left( c - \frac{t}{2} \right) \right) \\ &= -\frac{\cosh \frac{c}{2} \sinh \frac{s}{2}}{\cosh \frac{s}{2} \sinh \frac{c}{2}} \sinh s \cosh \frac{t}{2} \cosh \left( c - \frac{t}{2} \right) + \cosh s \cosh \frac{t}{2} \sinh \left( c - \frac{t}{2} \right) \\ &\quad - \sinh \frac{t}{2} \cosh \left( c - \frac{t}{2} \right), \end{aligned}$$

$$d|l_C| = \frac{\partial |l_C|}{\partial t} dt + \frac{\partial |l_C|}{\partial c} dc.$$

The two tangent vectors  $d|l_{DE}|$ ,  $d|l_C|$  are nonzero vectors. When  $c > 0$ ,  $0 \leq t \leq c$ ,

$$\frac{\partial |l_{DE}|}{\partial t} < 0, \quad \frac{\partial |l_C|}{\partial t} < 0.$$

For any  $k \leq 0$ ,  $d|l_{DE}| \neq kd|l_C|$ . Then there is a vector  $(A(c, t), B(c, t))$  such that

$$d|l_{DE}|(A(c, t), B(c, t)) > 0, \quad d|l_C|(A(c, t), B(c, t)) > 0.$$



By the assumption that  $l_{DE}$  lifts to a systole of  $X_0$ , only  $l_{DE}$  and  $l_C$  can lift to a systole of the surface. Then there is another surface with systole bigger than  $X_0$ . Therefore  $X_0$  is not maximal.  $\square$

From Propositions 7 and 8, we know that only  $l_{CE}$ ,  $l_{CD}$  and  $l_C$  in the orbifold  $S^2(2, 2, 2, n)$  can lift to a systole of the maximal surface.

By the symmetry of  $\Gamma(2, n)$  surfaces and Lemma 3, the preimage of the geodesic  $l_{CE} \subset S^2(2, 2, 2, n)$  ( $l_{CD}$ ,  $l_C$ , respectively) on the  $\Gamma(2, n)$  surface consists of pairwise disjoint geodesics with equal length.

**Proposition 9.** *On the maximal  $\Gamma(2, n)$  surface  $X_0$ , a simple closed geodesic is a systole if and only if it is lifted from  $l_{CE}$ ,  $l_{CD}$  or  $l_C$ .*

*Proof.* It is sufficient to prove that in  $X_0$ , every geodesic lifted from  $l_{CE}$ ,  $l_{CD}$  or  $l_C$  is a systole.

The proof is divided into two steps.

(1) If there is only one curve among  $l_{CE}$ ,  $l_{CD}$  and  $l_C$  that lifts to the systoles of  $X_0$ , then  $X_0$  is not maximal.


Without loss of generality, we assume that  $l_{CE}$  lifts to the systole of  $X_0$ , while  $l_{CD}$  and  $l_C$  do not lift to systoles of  $X_0$ . On the orbifold  $S^2(2, 2, 2, n)_{X_0}$ , there are deformations increasing or decreasing the length of the curve  $l_{CE}$ . A deformation increasing the length of  $l_{CE}$  increases the length of geodesics lifted from  $l_{CE}$  in  $X_0$ . If the deformation is small enough, then we get a new  $\Gamma(2, n)$  surface, whose systoles are lifted from  $l_{CE}$  and the length of these curves are longer than the corresponding curves in  $X_0$ . Hence  $X_0$  is not maximal.

(2) If there are exactly two curves among  $l_{CE}$ ,  $l_{CD}$  and  $l_C$  lifting to the systole of the  $\Gamma(2, n)$  surface, then the surface is not maximal.

(2a) We assume  $l_{CE}$  and  $l_{CD}$  lift to the systoles of  $X_0$ , while  $l_C$  does not lift to systoles of  $X_0$ . Then in the Fenchel–Nielsen coordinate  $(c, t)$ , the length  $|l_{CD}|(c, t)$  is monotonely increasing with respect to  $t$  by (5-3), while  $l_{CE} = c/2$ . We pick a sufficiently small  $\varepsilon > 0$  and deform the Fenchel–Nielsen coordinate from  $(c, t)$  to  $(c, t + \varepsilon)$ ; then we get a new surface  $X'$ . The systoles of  $X'$  are exactly the geodesics lifted from  $l_{CE}$ , and this surface has the same systole length to  $X_0$ . Then by (1), there exists a surface with longer systole than these two surfaces.

(2b) If  $l_{CE}$  and  $l_C$  lift to the systoles of  $X_0$ , while  $l_{CD}$  does not lift to systoles of  $X_0$ , the proof is similar. By (5-7),  $l_C$  is decreasing with respect to  $t$  when  $t \leq c$ . Thus using the deformation  $(c, t) \mapsto (c, t - \varepsilon)$ , we get a surface whose systoles are all lifted from  $l_{CE}$  and whose systole length is equal to  $X_0$ 's. Thus by (1), we know  $X_0$  is not maximal.

(2c) The last case is that  $l_{CD}$  and  $l_C$  lift to the systoles of  $X_0$ , while  $l_{CE}$  does not lift to systoles of  $X_0$ . Similarly, some of the Fenchel–Nielsen deformations along



**Figure 16.** Cutting off a subsurface with signature  $(1, 2)$ .

$l_{CD}$  increase the length of  $l_C$  and let  $l_{CD}$  become the unique geodesic that can lift to systoles of  $X_0$ . Then by (1), this surface is not maximal.

By (1) and (2), all three curves  $l_{CD}$ ,  $l_{CE}$  and  $l_C$  lift to the systoles of the maximal  $\Gamma(2, n)$  surface.  $\square$

We are now ready to calculate the maximal systole on hyperbolic  $\Gamma(2, n)$  surfaces.

*Proof of Theorem 1.* First we describe the lift of  $l_{CE}$ ,  $l_{CD}$  and  $l_C$  in the  $\Gamma(2, n)$  surface, respectively.


In the  $\Gamma(2, n)$  surface shown in Figure 16,  $C_i$  and  $C'_i$  ( $i = 1, \dots, n$ ) are the fixed points of the order-two involution  $\tau$  (recall its definition in Section 2) and are the lifts of the singular point  $C$  in  $S^2(2, 2, 2, n)$ ;  $D_i$  and  $D'_i$  are the mid-points of the seams, and the lifts of the point  $D$  in  $S^2(2, 2, 2, n)$ ;  $E_i$  and  $E'_i$  are the mid-points of  $C_i C'_i$  and the lifts of the point  $E$  in  $S^2(2, 2, 2, n)$ .

The curve  $l_{CE}$  lifts to cuffs of the surface, denoted as  $\gamma_i$  ( $i = 1, \dots, 5$ );  $l_{CD}$  lifts to geodesics passing through  $C_i D_i C'_{i+1} D'_i$  denoted as  $\alpha_i$ ;  $l_C$  lifts to geodesics passing through  $C_i C_{i+1}$ , denoted as  $\beta_i$ .

To calculate the systole length, we cut off a subsurface with signature  $(1, 2)$  from the  $\Gamma(2, n)$  surface containing  $\gamma_1$ ,  $\gamma_2$  and  $\alpha_1$  (Figures 16 and 17), the boundary length of this surface is given by [Buser 2010, p. 454, 2.4.1 (i)]. We take common perpendiculars between cuffs and boundary components as in Figure 17. Then in the hexagon  $H_1 H_2 A_1 A_2 A_3 A_4$ , we have

$$(5-8) \quad \begin{aligned} \cosh|H_1 H_2| &= \sinh|A_1 A_2| \sinh|A_3 A_4| \cosh|A_2 A_3| - \cosh|A_1 A_2| \cosh|A_3 A_4| \\ &= \sinh^2 c \cosh s - \cosh^2 c. \end{aligned}$$

In this subsurface,  $C_1, C'_1, C_2, C'_2$  are branched points of the hyperelliptic involution and  $\alpha_1, \beta_1, \gamma_1, \gamma_2$  are the systoles of the  $\Gamma(2, n)$  surface (Figure 18). To calculate the systole length, we redraw the  $(1, 2)$ -subsurface as Figure 19. When the systole



**Figure 17.** The red curves are cuffs. The blue curves are seams.

is maximal, the lengths of  $\gamma_1$  and  $\beta_1$  are the same, namely  $|C_1C'_1| = |C_1C_2|$ . In this case, the subsurface is shown in Figure 20.

When the surface is maximal,  $|H_5H_6| = \frac{1}{2}|\alpha_1| = c$ ,  $|C_1C'_1| = \frac{1}{2}\gamma_1 = c$ . We assume  $|H_3H_4| = l$  and  $|H_5C'_1| = h$ . Then in the hexagon  $H_4H_3C'_2H_6H_5C_1$ , by the symmetry of this hexagon,  $|H_6C'_2| = |H_5C'_1| = h$ , and by [Buser 2010, p. 454, 2.4.1(i)] we have


$$\cosh|H_3H_4| = \sinh|C_1H_5| \sinh|C'_2H_6| \cosh|H_5H_6| - \cosh|C_1H_5| \cosh|C'_2H_6|$$

and


$$(5-9) \quad \cosh l = \sinh^2 h \cosh c - \cosh^2 h.$$

In the triangle  $\triangle C_1H_5C'_1$ , by [Buser 2010, p. 454, 2.2.2 (i)], we have

$$(5-10) \quad \cosh|C_1C'_1| = \cosh|C_1H_5| \cosh|C'_1H_5| \cosh c = \cosh h \cosh \frac{c}{2}.$$



**Figure 18.** Systoles of  $X$  in the  $(1, 2)$ -subsurface.



**Figure 19.** Redrawing the  $(1, 2)$ -subsurface.

For convenience, we denote  $\cosh c$  by  $K$ . Then combining (5-9) and (5-10), we eliminate  $h$  and get

$$(5-11) \quad \frac{2K^2}{K+1} = \frac{K + \cosh l}{K-1}.$$

Recall that  $l = |H_3 H_4|$ . Then combining (5-11), (5-8) and (5-1), we eliminate  $l$  and  $s$ , and get

$$2K^3 - 3K^2 + 1 - 4 \cos^2 \frac{\pi}{n} (K+1)^2 = 0.$$

The unique real solution of this equation is


$$K = \sqrt[3]{\frac{1}{216}L^3 + \frac{1}{8}L^2 + \frac{5}{8}L - \frac{1}{8} + \sqrt{\frac{1}{108}L(L^2 + 18L + 27)}} + \sqrt[3]{\frac{1}{216}L^3 + \frac{1}{8}L^2 + \frac{5}{8}L - \frac{1}{8} - \sqrt{\frac{1}{108}L(L^2 + 18L + 27)}} + \frac{1}{6}(L + 3),$$

where  $L = 4 \cos^2 \frac{\pi}{n}$ .

At last, we calculate the twist parameter  $t$  of the maximal surface, using (5-1) and (5-3):

$$\cosh \frac{t}{2} = \frac{\cosh \frac{c}{2}}{\cosh \frac{s}{2}} = \frac{\cosh^2 \frac{c}{2}}{\cos \frac{\pi}{n}} = \frac{\cosh c + 1}{2 \cos \frac{\pi}{n}} = \frac{K + 1}{2 \cos \frac{\pi}{n}}.$$

Theorem 1 follows. □



**Figure 20.** When  $|C_1 C'_1| = |C_1 C_2|$ .

### Acknowledgements

We would like to thank Professors Sheng Bai and Shicheng Wang for their helpful discussions and suggestions. Gao thanks Professor Shicheng Wang for his supervision, help and support during his PhD studies. We thank Professor Ying Zhang for his helpful comments.

Wang is partially supported by NSFC 12131009 and National Key R&D Program of China 2020YFA0712800.

### References

- [Bai et al. 2021] S. Bai, Y. Gao, and S. Wang, “Systoles of hyperbolic surfaces with big cyclic symmetry”, *Sci. China Math.* **64**:2 (2021), 421–442. MR Zbl
- [Ballmann et al. 2016] W. Ballmann, H. Matthiesen, and S. Mondal, “Small eigenvalues of closed surfaces”, *J. Differential Geom.* **103**:1 (2016), 1–13. MR Zbl
- [Ballmann et al. 2018] W. Ballmann, H. Matthiesen, and S. Mondal, “Small eigenvalues of surfaces: old and new”, *ICCM Not.* **6**:2 (2018), 9–24. MR Zbl
- [Bavard 1992] C. Bavard, “La systole des surfaces hyperelliptiques”, preprint 71, Ec. Norm. Sup. Lyon, 1992.
- [Buser 2010] P. Buser, *Geometry and spectra of compact Riemann surfaces*, Birkhäuser, Boston, 2010. MR Zbl
- [Buser and Sarnak 1994] P. Buser and P. Sarnak, “On the period matrix of a Riemann surface of large genus”, *Invent. Math.* **117**:1 (1994), 27–56. MR Zbl
- [Chen and Li 2015] L. Chen and W. Li, “Systoles of surfaces and 3-manifolds”, pp. 61–80 in *Geometry and topology of submanifolds and currents*, edited by W. Li and S. W. Wei, Contemp. Math. **646**, Amer. Math. Soc., Providence, RI, 2015. MR Zbl
- [Croke and Katz 2003] C. B. Croke and M. Katz, “Universal volume bounds in Riemannian manifolds”, pp. 109–137 in *Surveys in differential geometry* (Boston, 2002), vol. 8, edited by S.-T. Yau, Surv. Differ. Geom. **8**, International Press, Somerville, MA, 2003. MR Zbl
- [Fortier Bourque and Rafi 2022] M. Fortier Bourque and K. Rafi, “Local maxima of the systole function”, *J. Eur. Math. Soc. (JEMS)* **24**:2 (2022), 623–668. MR Zbl
- [Gromov 1983] M. Gromov, “Filling Riemannian manifolds”, *J. Differential Geom.* **18**:1 (1983), 1–147. MR Zbl
- [Jenni 1984] F. Jenni, “Über den ersten Eigenwert des Laplace-operators auf ausgewählten Beispielen kompakter Riemannscher Flächen”, *Comment. Math. Helv.* **59**:2 (1984), 193–203. MR Zbl
- [Katz et al. 2007] M. G. Katz, M. Schaps, and U. Vishne, “Logarithmic growth of systole of arithmetic Riemann surfaces along congruence subgroups”, *J. Differential Geom.* **76**:3 (2007), 399–422. MR Zbl
- [Mondal 2014] S. Mondal, “Systole and  $\lambda_{2g-2}$  of closed hyperbolic surfaces of genus  $g$ ”, *Enseign. Math.* **60**:1-2 (2014), 3–24. MR Zbl
- [Mumford 1971] D. Mumford, “A remark on Mahler’s compactness theorem”, *Proc. Amer. Math. Soc.* **28** (1971), 289–294. MR Zbl
- [Parlier 2014] H. Parlier, “Simple closed geodesics and the study of Teichmüller spaces”, pp. 113–134 in *Handbook of Teichmüller theory, IV*, edited by A. Papadopoulos, IRMA Lect. Math. Theor. Phys. **19**, Eur. Math. Soc., Zürich, 2014. MR Zbl

- [Petri 2018] B. Petri, “Hyperbolic surfaces with long systoles that form a pants decomposition”, *Proc. Amer. Math. Soc.* **146**:3 (2018), 1069–1081. MR Zbl
- [Petri and Walker 2018] B. Petri and A. Walker, “Graphs of large girth and surfaces of large systole”, *Math. Res. Lett.* **25**:6 (2018), 1937–1956. MR
- [Schmutz 1993] P. Schmutz, “Riemann surfaces with shortest geodesic of maximal length”, *Geom. Funct. Anal.* **3**:6 (1993), 564–631. MR Zbl
- [Wang et al. 2013] C. Wang, S. Wang, Y. Zhang, and B. Zimmermann, “Extending finite group actions on surfaces over  $S^3$ ”, *Topology Appl.* **160**:16 (2013), 2088–2103. MR Zbl
- [Wang et al. 2015] C. Wang, S. Wang, Y. Zhang, and B. Zimmermann, “Embedding surfaces into  $S^3$  with maximum symmetry”, *Groups Geom. Dyn.* **9**:4 (2015), 1001–1045. MR Zbl
- [Wolpert 2017] S. A. Wolpert, “Equiboundedness of the Weil–Petersson metric”, *Trans. Amer. Math. Soc.* **369**:8 (2017), 5871–5887. MR Zbl
- [Wu 2019] Y. Wu, “Growth of the Weil–Petersson inradius of moduli space”, *Ann. Inst. Fourier (Grenoble)* **69**:3 (2019), 1309–1346. MR Zbl

Received November 28, 2021. Revised June 18, 2023.

YUE GAO  
SCHOOL OF MATHEMATICS AND STATISTICS  
ANHUI NORMAL UNIVERSITY  
WUHU  
CHINA  
yuegao@ahnu.edu.cn

JIAJUN WANG  
LMAM, SCHOOL OF MATHEMATICAL SCIENCES  
PEKING UNIVERSITY  
BEIJING  
CHINA  
wjiajun@pku.edu.cn

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 325    No. 1    July 2023

---

Estimate for the first fourth Steklov eigenvalue of a minimal hypersurface with free boundary	1
RONDINELLE BATISTA, BARNABÉ LIMA, PAULO SOUSA and BRUNO VIEIRA	
Catenoid limits of singly periodic minimal surfaces with Scherk-type ends	11
HAO CHEN, PETER CONNOR and KEVIN LI	
The strong homotopy structure of BRST reduction	47
CHIARA ESPOSITO, ANDREAS KRAFT and JONAS SCHNITZER	
The maximal systole of hyperbolic surfaces with maximal $S^3$ -extendable abelian symmetry	85
YUE GAO and JIAJUN WANG	
Stable systoles of higher rank in Riemannian manifolds	105
JAMES J. HEBDA	
Spin Kostka polynomials and vertex operators	127
NAIHUAN JING and NING LIU	
The structure of groups with all proper quotients virtually nilpotent	147
BENJAMIN KLOPSCH and MARTYN QUICK	